

# Testing Mediators of Intervention Effects in Randomized Controlled Trials: An Evaluation of Three Depression Prevention Programs

Eric Stice, Paul Rohde, John R. Seeley, and Jeff M. Gau  
Oregon Research Institute

**Objective:** Evaluate a new 5-step method for testing mediators hypothesized to account for the effects of depression prevention programs. **Method:** In this indicated prevention trial, at-risk teens with elevated depressive symptoms were randomized to a group cognitive-behavioral (CB) intervention, group supportive expressive intervention, CB bibliotherapy, or assessment-only control condition. **Results:** The group CB intervention reduced depressive symptoms and negative cognitions and increased pleasant activities. Change in these mediators predicted change in depression, and intervention effects became weaker controlling for change in the mediators; yet, change in depression appeared typically to occur before change in the mediators. The supportive expressive intervention reduced depressive symptoms but affected only 1 of 2 mediators (emotional expression but not loneliness). Change in emotional expression did not correlate with change in depression, and change in depression usually occurred before change in the mediators. Bibliotherapy did not significantly affect depressive symptoms or the ostensive mediators (negative cognitions and pleasant activities), and change in depression usually occurred before change in the mediators. **Conclusion:** Results imply that this procedure provides a sensitive test of mediation but yielded limited support for the hypothesized mediators, suggesting that nonspecific factors may play an important mediational role.

**Keywords:** mediation, depression, prevention, adolescents

Cognitive-behavioral (CB) depression prevention programs, which focus on reducing negative cognitions and increasing pleasant activities, have reduced depressive symptoms and risk for future onset of major depression relative to assessment-only control conditions in randomized trials (Stice, Shaw, Bohon, Marti, & Rohde, 2009). Yet, few trials have investigated mediators that account for these effects. Such an investigation provides a test of the theoretical mechanisms that putatively underlie program effects and can implicate components that are most effective, potentially aiding in program refinement (Kazdin & Nock, 2003).

Although procedures for examining mediators that account for intervention effects have been proposed, they do not provide a rigorous test of all of the components necessary for mediation, particularly the temporal sequencing between changes in the mediator and the outcome. As has been noted, it is vital to document that change in the mediator temporally precedes change in the outcome (Kazdin & Nock, 2003; Weersing & Weisz, 2002). Because a specific procedure for testing this temporal sequencing of

change has not been offered, we proposed a five-step procedure for testing mediation with data from randomized trials (Stice, Presnell, Gau, & Shaw, 2007) and suggested that the strongest case for mediation would be made when five conditions are satisfied: (a) intervention participants show greater decreases on the outcome than do controls; (b) intervention participants show greater decreases on the mediator than do controls; (c) change in the mediator predicts change in the outcome in the intervention condition;<sup>1</sup> (d) the predictive effect of the intervention on change in the outcome is significantly reduced (for partial mediation) or eliminated (for full mediation), controlling for change in the mediator; and (e) meaningful change in the mediator occurs before meaningful change in the outcome more frequently than would be expected by chance in the intervention condition. We think this five-step procedure provides a rigorous approach for testing mediation in randomized trials, yet this heuristic model may require refinement (e.g., the fifth criterion offers but one method for testing whether the mediator changes before the outcome) and these criteria involve assumptions that should be verified.

Although trials have evaluated mediators that account for the effects of depression prevention programs, the most rigorous ones have examined only Criteria 1 through 4 (Jaycox, Reivich, Gillham, & Seligman, 1994; Yu & Seligman, 2002), with mixed results. Our

---

Eric Stice, Paul Rohde, John R. Seeley, and Jeff M. Gau, Oregon Research Institute.

This study was supported by National Institute of Health Grants MH067183 and MH080760. Our thanks go to project research assistants Courtney Byrd, Kathryn Fischer, Amy Folmer, Cassie Goodin, Jacob Mase, and Emily Wade; a multitude of undergraduate volunteers; the Austin Independent School District; and the participants who made this study possible.

Correspondence concerning this article should be addressed to Eric Stice, Oregon Research Institute, 1715 Franklin Boulevard, Eugene, OR 97403. E-mail: estice@ori.org

---

<sup>1</sup> We acknowledge that change in the mediator may predict change in the outcome in the control condition. Yet, we assert that, because the change in the mediator in the control condition could not be caused by the intervention, the correlation between the mediator and outcome among controls reflects etiologic or maintenance processes that are not germane to testing mediators that account for intervention effects.

purpose in the present report was to evaluate the five-step method for testing the hypothesized mediators of depression prevention programs. In this indicated depression prevention program, at-risk youths with elevated depressive symptoms were randomized to a CB group, a supportive expressive group, CB bibliotherapy, or an assessment-only control condition. We hypothesized that reductions in negative cognitions and increases in pleasant activities would mediate the effects of the group and bibliotherapy CB programs and that increases in emotional expression and reductions in loneliness would mediate the effects of the supportive expressive program on reductions in depressive symptoms. We tested for mediation during intervention delivery, because this is typically when the most pronounced intervention effects occur; also, change is often linear during this period (Stice et al., 2009), making it easier to model change. As suggested by Kazdin and Nock (2003), changes were assessed in the mediator and outcome during the 6-week intervention delivery at baseline, midway through, and after termination. Although we considered using weekly surveys, participants complained about the assessment burden in our prior trial that took this approach (Stice et al., 2007). A trial evaluating multiple programs also affords an opportunity to examine the specificity of intervention effects. Kazdin and Nock (2003) argued that it is important to test whether interventions affect mediators specific to that intervention but not mediators specific to other interventions. Thus, we also tested whether each prevention program affects the theoretically specific mediators but not mediators specific to the other programs.

### Method

Participants were 341 high school students (mean age = 15.6 years,  $SD = 1.2$ ) who reported elevated depressive symptoms. Participants were 2% Asian, 9% African American, 46% Caucasian, 33% Hispanic, and 10% with mixed heritage. Their ethnic distribution was more diverse than that of the population. Details regarding recruitment, intervention content, supervision, therapist fidelity and competence, and attrition are provided by Stice, Rohde, Seeley, and Gau (2008).

*Depressive symptoms* were assessed with the 21-item Beck Depression Inventory (BDI; Beck, Steer, & Garbin, 1988). The BDI has shown internal consistency ( $\alpha = .73-.95$ ), test-retest reliability ( $r_s = .60-.90$ ), and convergent validity with clinician ratings of depressive symptoms in adults (mean  $r = .75$ ; Beck et al., 1988). The BDI showed internal consistency ( $\alpha = .89$ ) and 3-week test-retest reliability in the control condition ( $r = .76$ ), and it correlated with functional impairment ( $r = .54$ ); assessed by the Social Adjustment Scale-Self Report for Youth; Weissman, Orvaschel, & Padian, 1980) and self-esteem ( $r = -.53$ ; assessed with the scale from Rosenberg, 1979).

*Negative cognitions* were assessed with 12 items from the Automatic Thoughts Questionnaire (ATQ; Hollon & Kendall, 1980). The original 30-item ATQ scale correlated more strongly with depressive than with anxiety symptoms in adolescent inpatients (Jolly & Wiesner, 1996). The 12 items from the personal maladjustment/desire for change and negative self-concept/expectations subscales were used because they correlated .98 with the total ATQ score in a past adolescent depression treatment trial (Rohde, Clarke, Mace, Jorgensen, & Seeley, 2004).<sup>2</sup> Pilot testing with teens for this trial ( $N = 44$ ) indicated that a 12-item short

form of the ATQ showed internal consistency ( $\alpha = .92$ ) and 1-week test-retest reliability ( $r = .74$ ) and that it correlated with BDI scores ( $r = .62$ ). The 12-item ATQ showed internal consistency ( $\alpha = .93$ ) and 3-week test-retest reliability ( $r = .75$ ), and it correlated with BDI scores ( $r = .65$ ), functional impairment ( $r = .53$ ), and self-esteem ( $r = -.59$ ).

*Pleasant activities* were assessed with 12 items from the Pleasant Events Schedule (PES; MacPhillamy & Lewinsohn, 1982) commonly endorsed by teens. The original 320-item PES showed 72% agreement with peer ratings of pleasant activities, had predictive validity for future self-reported pleasant activities (mean  $r = .59$ ), and discriminated between depressed and healthy adults (MacPhillamy & Lewinsohn, 1982). This 12-item scale correlated with the total PES score ( $r = .84$ ) in an adolescent depression treatment trial (Rohde et al., 2004). Pilot testing with teens ( $N = 44$ ) indicated that the 12-item form of the PES showed internal consistency ( $\alpha = .79$ ), had 1-week test-retest reliability ( $r = .88$ ), and correlated with BDI scores ( $r = -.45$ ). This scale showed internal consistency ( $\alpha = .73$ ) and 3-week test-retest reliability ( $r = .62$ ), and it correlated with BDI scores ( $r = -.37$ ), functional impairment ( $r = -.36$ ), and self-esteem ( $r = .36$ ).

*Emotional expression* was assessed with nine items developed for this study (e.g., "I was able to discuss my feelings in a helpful way"), because we were unable to locate a scale developed for teens that assessed this construct. This scale showed internal consistency ( $\alpha = .82$ ) and 3-week test-retest reliability ( $r = .66$ ), and it correlated with BDI scores ( $r = -.29$ ), functional impairment ( $r = -.29$ ), and self-esteem ( $r = .39$ ).

*Loneliness* was assessed with an eight-item Loneliness Scale adapted by Lewinsohn et al. (1994) from Russell (1996). The full 20-item Loneliness Scale correlated with other loneliness scales (mean  $r = .69$ ) and correlated inversely with self-reported social support (mean  $r = -.52$ ) among adults (Russell, 1996). The eight-item scale had a correlation of .92 with the full 20-item scale in a second pilot study of teens ( $n = 25$ ). It showed internal consistency ( $\alpha = .83$ ) and 3-week test-retest reliability ( $r = .74$ ), and it correlated with BDI scores ( $r = .49$ ), functional impairment ( $r = .53$ ), and self-esteem ( $r = -.51$ ).

### Results

Random coefficient unconditional growth models confirmed the a priori assumption that linear trends were appropriate for modeling change over time for the mediators and outcome (see Table 1 for cell means across assessment points). Stice et al. (2007) reported details regarding the analytic approach. Hierarchical linear

<sup>2</sup> We report the correlations between the short forms of the negative cognitions, pleasant activities, and loneliness scales used herein and the original full-length forms in the Measures section, because these correlations provide an indication of how well the short forms serve as a proxy for the longer forms. In this footnote, we report the correlations between the short forms of these scales and the items from the full-length forms not included in the short forms of these scales, which is akin to the split-half reliability of these scales, as this may be of interest to some readers (negative cognitions,  $r = .96$ ; pleasant activities,  $r = .78$ ; loneliness,  $r = .76$ ).

Table 1  
Means and Standard Deviations for the Raw Outcome and Mediator Variables at Each of the Assessments

Mediator, outcome, and assessment	CBT ( <i>n</i> = 89)		Supportive ( <i>n</i> = 88)		Bibliotherapy ( <i>n</i> = 80)		Control ( <i>n</i> = 84)	
	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>
Beck Depression Inventory								
Time 1	20.00	10.35	20.27	9.83	18.20	7.53	19.60	9.23
Time 2	15.28	8.19	16.87	9.33	16.74	9.09	18.92	9.33
Time 3	10.78	9.04	14.68	10.53	14.51	8.97	16.80	9.76
Negative cognitions								
Time 1	2.60	0.94	2.56	0.95	2.42	0.82	2.47	0.79
Time 2	2.30	0.74	2.29	0.93	2.29	0.78	2.45	0.85
Time 3	1.95	0.77	2.18	1.01	2.13	0.84	2.28	0.83
Pleasant activities								
Time 1	23.67	9.31	24.28	9.32	24.94	7.40	24.04	9.59
Time 2	25.17	8.22	26.96	8.28	24.30	8.26	25.36	7.89
Time 3	29.01	9.44	28.42	9.28	25.24	9.18	25.55	8.59
Emotional expression								
Time 1	2.73	0.82	3.00	0.76	2.92	0.74	2.91	0.67
Time 2	3.12	0.77	3.32	0.80	2.96	0.65	2.95	0.68
Time 3	3.48	0.93	3.62	0.77	3.11	0.84	3.04	0.80
Loneliness								
Time 1	23.57	5.99	21.61	7.00	22.12	6.29	22.25	5.97
Time 2	20.93	5.86	19.89	5.56	21.21	5.32	21.81	5.99
Time 3	18.99	5.93	20.12	5.87	20.83	6.40	21.40	6.22

Note. BDI = Beck Depression Inventory; CBT = cognitive-behavioral therapy.

modeling was used because it produces empirical Bayesian estimates, which were used in our procedure for testing mediation.<sup>3</sup>

### Criterion 1

While controlling for the depression intercept, we tested whether intervention condition predicted the slope parameter for depressive symptoms. CB group (see Table 2) and supportive expressive participants (see Table 4), but not bibliotherapy participants (see Table 3), showed significantly greater reductions in depression than did assessment-only controls.

### Criterion 2

We tested whether intervention condition predicted the slope parameter for each mediator, controlling for the intercept of the respective mediator. CB group participants (see Table 2) but not bibliotherapy participants (see Table 3) showed significantly greater decreases in negative cognitions and greater increases in pleasant activities than did controls (see Table 2). Supportive expressive participants showed significantly greater increases in emotional expression relative to controls, but there were no effects for loneliness (see Table 4).

We also tested whether the intervention effects on the mediators were specific. Relative to controls, group CB participants showed greater decreases in loneliness ( $B = -1.86$ ,  $SE = 0.44$ ,  $t = -4.22$ ,  $p < .001$ ) and increases in emotional expression ( $B = 0.31$ ,  $SE = 0.06$ ,  $t = 5.00$ ,  $p < .001$ ). Bibliotherapy participants did not show differential change in loneliness ( $B = -0.22$ ,  $SE = 0.40$ ,  $t = -0.57$ ,  $p = .572$ ) or in emotional expression ( $B = 0.03$ ,  $SE = 0.06$ ,  $t = 0.47$ ,  $p = .637$ ). Supportive expressive participants showed greater increases in pleasant activities ( $B = 1.31$ ,  $SE =$

$0.60$ ,  $t = 2.18$ ,  $p = .030$ ) but not in negative cognitions ( $B = -0.09$ ,  $SE = 0.06$ ,  $t = -1.53$ ,  $p = .128$ ).

### Criterion 3

We tested whether the slope of the mediator predicted change in depressive symptoms, controlling for the depression intercept. Change in negative cognitions and pleasant activities predicted change in depressive symptoms for the CB group (see Table 2) and bibliotherapy condition (see Table 3). Change in loneliness, but not in emotional expression, predicted change in depressive symptoms for those in the supportive expressive condition (see Table 4).

<sup>3</sup> We noted errors in the equations provided in Stice et al. (2007). Each of the composite multilevel models has the time parameter omitted from the rate-of-change random effect; the person-level random effect is represented with the Greek letter zeta rather than the more standard  $e$ ; and the composite model for Criterion 4 incorrectly identifies  $\gamma_{12}$  as a group by change in mediator parameter. Important to note is the fact that errors were only in the description of the equations; the models used to generate the parameters estimates reported in the article are correct. In particular, on p. 24, column 2, the equation on lines 7–8 should read  $Y_{ij} = [\gamma_{00} + \gamma_{10}(\text{time}_{ij}) + \gamma_{01}(\text{group}_i) + \gamma_{11}(\text{Group}_i \times \text{Time}_{ij})] + [\zeta_{0i} + \zeta_{1i}(\text{time}_{ij}) + e_{ij}]$ ; the parenthetical ( $\zeta_{ij}$ ) on line 15 should read ( $e_{ij}$ ); the equation on lines 18–20 should read  $Y_{ij} = [\gamma_{00} + \gamma_{10}(\text{time}_{ij}) + \gamma_{01}(\Delta\text{mediator}_i) + \gamma_{11}(\Delta\text{Mediator}_i \times \text{Time}_{ij})] + [\zeta_{0i} + \zeta_{1i}(\text{time}_{ij}) + e_{ij}]$ ; the parenthetical ( $\zeta_{ij}$ ) on line 27 should read ( $e_{ij}$ ); the equation on lines 30–32 should read  $Y_{ij} = [\gamma_{00} + \gamma_{10}(\text{time}_{ij}) \pm \gamma_{01}(\text{group}_i) + \gamma_{02}(\Delta\text{mediator}_i) \pm \gamma_{11}(\text{Group}_i \times \text{Time}_{ij})] + \gamma_{12}(\Delta\text{Mediator}_i \times \text{Time}_{ij}) + [\zeta_{0i} + \zeta_{1i}(\text{time}_{ij}) + e_{ij}]$ ; the phrase “effect of condition on the change in the mediator” on lines 36–37 should read “effect of change in the mediator on change in the outcome”; and the parenthetical ( $\zeta_{ij}$ ) on line 41 should read ( $e_{ij}$ ).

Table 2  
*Test of Hypothesized Mediators of the Group CB Intervention Effects Compared to the Control Group*

Criterion	Fixed effects					Random effects	
	<i>B</i>	<i>SE</i>	<i>t</i> ratio	<i>p</i>	Pr	Between subjects	Within subjects
1. Effect of the treatment on the outcome; $\gamma_{II}$ CBT $\rightarrow$ $\Delta$ BDI	-3.21	0.72	-4.48	<.001	-.32	84.60	21.06
2. Effect of the treatment on the mediator; $\gamma_{II}$ CBT $\rightarrow$ $\Delta$ negative cognitions	-0.23	0.06	-4.07	<.001	-.30	0.63	0.17
CBT $\rightarrow$ $\Delta$ pleasant activities	1.91	0.68	2.82	.006	.21	66.81	22.48
3. Relation between change in mediator on change in the outcome; $\gamma_{II}$ $\Delta$ negative cognitions $\rightarrow$ $\Delta$ BDI	17.88	3.67	4.87	<.001	.46	81.12	19.29
$\Delta$ pleasant activities $\rightarrow$ $\Delta$ BDI	-1.01	0.26	-3.88	<.001	-.38	89.31	19.29
4a. Effect of treatment on outcome controlling for change in mediator; $\gamma_{II}$ CBT $\rightarrow$ $\Delta$ BDI   negative cognitions	0.53	0.85	0.62	.536	.05	68.23	21.06
CBT $\rightarrow$ $\Delta$ BDI   pleasant activities	-1.49	0.75	-2.00	.046	-.15	76.62	21.06
						<i>t</i> ratio	$\Delta$ Pr (%)
4b. Effect of treatment on outcome, controlling for change in mediator significantly reduced or eliminated compared to effect of the treatment on the outcome CBT $\rightarrow$ $\Delta$ BDI   negative cognitions						-6.73	116
CBT $\rightarrow$ $\Delta$ BDI   pleasant activities						-5.21	56
						% showing	<i>p</i>
5a. % showing .33 <i>SD</i> decrease in mediator before .33 <i>SD</i> in outcome for the CBT condition Decrease in negative cognitions before decrease in BDI						8.0	<.001
Increase in pleasant activities before decrease in BDI						9.0	<.001
5b. % showing .33 <i>SD</i> decrease in mediator before .33 <i>SD</i> in outcome for the control condition Decrease in negative cognitions before decrease in BDI						3.6	<.001
Increase in pleasant activities before decrease in BDI						8.3	<.001

Note. CB = cognitive-behavioral; *B* = unstandardized regression coefficient; *SE* = standard error; Pr = partial regression coefficient; CBT = cognitive-behavioral therapy;  $\Delta$  = change; BDI = Beck Depression Inventory; | = controlling for; *SD* = standard deviation.

#### Criterion 4

We tested whether the predictive effect of treatment on change in depressive symptoms was significant when controlling for change in the mediator (4a) and whether the predictive effect of treatment on change in depression was significantly reduced when controlling for change in the mediator (4b). For the CB group (see Table 2), treatment no longer predicted change in depression when controlling for change in negative cognitions, but treatment predicted change in depression when controlling for change in pleasant activities (4a). The effect of the CB group intervention on depression was significantly reduced when change in each of these mediators was controlled (4b). For bibliotherapy (see Table 3), treatment showed a significant relation to change in depression when change in negative cognitions and change in pleasant activities were controlled (4a), and there was a significant change in the effect of treatment on change in depression when change in the mediators was controlled (4b). These results appear to provide support for Criterion 4 for bibliotherapy; however, there was no main effect of bibliotherapy on change in depression, indicating that these results do not provide support for Criterion 4. The effect of the supportive expressive intervention on depression became nonsignificant when change in loneliness and emotional expression were controlled (4a). Yet, the effect of treatment on change in depression was significantly weaker only when change in loneliness was controlled (see Table 4).

#### Criterion 5

We tested whether meaningful reduction in the mediator typically occurred before meaningful reduction in the outcome more frequently than expected on the basis of chance. We defined meaningful change as a .33 *SD* reduction in the variables, because most participants showed this degree of change in the mediators and outcomes. Although this definition is somewhat arbitrary, the important point is that we used the same criterion for mediators and for the outcome that adjust for the fact that the measures had different scaling. Participants who showed a .33 *SD* reduction in the mediator before they showed a .33 *SD* reduction in the outcome received a score of 1. Those who showed a .33 *SD* decrease in the mediator after they showed a .33 *SD* decrease in the outcome, did not show a .33 *SD* decrease in the outcome, or did not show a .33 *SD* decrease in the mediator received a score of 0. We used a nonparametric binomial test to determine whether the proportion of participants who showed a meaningful change in the mediator before showing a meaningful change in the outcome was greater than .50 among intervention participants. In the CB group condition, 75% showed a .33 *SD* decrease in depression, but only 8% showed a decrease in negative cognitions before a .33 *SD* reduction in depression and only 9% showed an increase in pleasant activities before a .33 *SD* reduction in depression (see Table 2). In the bibliotherapy condition, 55% showed a .33

Table 3  
*Test of Hypothesized Mediators of the Bibliotherapy Intervention Effects Compared to the Control Group*

Criterion	Fixed effects					Random effects	
	<i>B</i>	<i>SE</i>	<i>t</i> ratio	<i>p</i>	Pr	Between subjects	Within subjects
1. Effect of the treatment on the outcome; $\gamma_{11}$ Bibliotherapy $\rightarrow$ $\Delta$ BDI	-0.45	0.57	-0.80	.430	-.06	58.86	20.12
2. Effect of the treatment on the mediator; $\gamma_{11}$ Bibliotherapy $\rightarrow$ $\Delta$ negative cognitions	-0.05	0.05	-0.98	.328	-.08	0.50	0.18
Bibliotherapy $\rightarrow$ $\Delta$ pleasant activities	-0.61	0.61	-1.00	.318	-.08	53.05	20.23
3. Relation between change in mediator on change in the outcome; $\gamma_{11}$ $\Delta$ negative cognitions $\rightarrow$ $\Delta$ BDI	50.01	8.21	6.09	<.001	.57	49.76	17.16
$\Delta$ pleasant activities $\rightarrow$ $\Delta$ BDI	-1.14	0.28	-4.05	<.001	-.42	52.37	17.16
4a. Effect of treatment on outcome controlling for change in mediator; $\gamma_{11}$ Bibliotherapy $\rightarrow$ $\Delta$ BDI   negative cognitions	1.75	0.57	3.10	.003	.24	56.99	19.47
Bibliotherapy $\rightarrow$ $\Delta$ BDI   pleasant activities	-1.07	0.54	-1.97	.050	-.15	57.48	20.12
						<i>t</i> ratio	$\Delta$ Pr (%)
4b. Effect of treatment on outcome, controlling for change in mediator significantly reduced or eliminated compared to effect of the treatment on the outcome Bibliotherapy $\rightarrow$ $\Delta$ BDI   negative cognitions						-7.68	400
Bibliotherapy $\rightarrow$ $\Delta$ BDI   pleasant activities						4.92	-150
						% showing	<i>p</i>
5a. % showing .33 <i>SD</i> decrease in mediator before .33 <i>SD</i> in outcome for the CBT condition							
Decrease in negative cognitions before decrease in BDI						6.3	<.001
Increase in pleasant activities before decrease in BDI						5.0	<.001
5b. % showing .33 <i>SD</i> decrease in mediator before .33 <i>SD</i> in outcome for the control condition							
Decrease in negative cognitions before decrease in BDI						1.2	<.001
Increase in pleasant activities before decrease in BDI						10.7	<.001

Note. *B* = unstandardized regression coefficient; *SE* = standard error; Pr = partial regression coefficient;  $\Delta$  = change; BDI = Beck Depression Inventory; | = controlling for; *SD* = standard deviation.

*SD* decrease in depression, but only 6% showed a decrease in negative cognitions before a .33 *SD* in reduction of depression and only 5% showed an increase in pleasant activities before a .33 *SD* reduction in depression (see Table 3). In the supportive expressive condition, 69% showed a .33 *SD* decrease in depression, but only 15% showed a decrease in loneliness before a .33 *SD* reduction in depression and only 39% showed an increase in emotional expression before a .33 *SD* reduction in depression (see Table 4). None of these proportions were significantly greater than .50. For reference, 65% of controls showed a .33 *SD* decrease in depression; between 1% and 16% of controls showed a .33 *SD* decrease in mediators before showing a .33 *SD* decrease in depression.

## Discussion

Results provided partial support for the intervention theory for the group CB. Participants showed greater reductions in depressive symptoms and greater change in negative cognitions and pleasant activities than did controls, with effect sizes that were clinically significant, and change in the mediators correlated with change in depression. The intervention effect became nonsignificant when change in negative cognitions was controlled but remained significant when change in pleasant activities was controlled, with the effects of the intervention on

change in depression being significantly reduced. Yet results suggested that only 8% of participants showed meaningful change in the mediators before showing meaningful change in depression; this was significantly lower than expected by chance. Thus, results do not appear to support the theory that changes in negative cognitions and pleasant activities mediate the intervention effects, because the outcome typically changed before the mediators. Given that the most rigorous tests of mediators of CB depression prevention programs examined only the first four criteria (Jaycox et al., 1994; Yu & Seligman, 2002), it is possible that change in depression may have preceded change in the putative mediator in those trials, too.

Results provided little support for the intervention theory for bibliotherapy. Relative to controls, bibliotherapy participants did not show greater improvements in depressive symptoms, negative cognitions, or pleasant activities. Although change in the mediators correlated with change in depression, there was no main effect of intervention on depression, so it was not relevant to test whether the main effects of treatment on the outcome became weaker when change in the mediator was controlled. Further, results imply that only 6% of participants showed a decrease in negative cognitions before showing a reduction in depression and that only 5% of participants showed an increase in pleasant activities before showing a reduction in

Table 4  
*Test of Hypothesized Mediators of the Supportive Intervention Effects Compared to the Control Group*

Criterion	Fixed effects					Random effects	
	<i>B</i>	<i>SE</i>	<i>t</i> ratio	<i>p</i>	Pr	Between subjects	Within subjects
1. Effect of the treatment on the outcome; $\gamma_{11}$ Supportive $\rightarrow$ $\Delta$ BDI	-1.40	0.62	-2.23	.027	-.17	74.02	22.22
2. Effect of the treatment on the mediator; $\gamma_{11}$ Supportive $\rightarrow$ $\Delta$ loneliness	-0.32	0.47	-0.69	.494	-.05	39.05	8.26
Supportive $\rightarrow$ $\Delta$ emotional expression	0.24	0.06	4.17	<.001	.30	0.36	0.19
3. Relation between change in mediator on change in the outcome; $\gamma_{11}$ $\Delta$ loneliness $\rightarrow$ $\Delta$ BDI	0.80	0.25	3.18	.002	.32	74.14	21.53
$\Delta$ emotional expression $\rightarrow$ $\Delta$ BDI	1.62	3.78	0.43	.688	.05	83.10	21.53
4a. Effect of treatment on outcome controlling for change in mediator; $\gamma_{11}$ Supportive $\rightarrow$ $\Delta$ BDI   loneliness	-1.11	0.58	-1.90	.059	-.14	67.96	22.22
Supportive $\rightarrow$ $\Delta$ BDI   emotional expression	-1.03	0.65	-1.58	.116	-.12	74.29	22.22
						<i>t</i> ratio	$\Delta$ Pr (%)
4b. Effect of treatment on outcome, controlling for change in mediator significantly reduced or eliminated compared to effect of the treatment on the outcome Supportive $\rightarrow$ $\Delta$ BDI   loneliness						-4.48	18
Supportive $\rightarrow$ $\Delta$ BDI   emotional expression						-1.86	29
						% showing	<i>p</i>
5a. % showing .33 <i>SD</i> decrease in mediator before .33 <i>SD</i> in outcome for the supportive condition Decrease in loneliness before decrease in BDI						14.8	<.001
Increase in emotional expression before decrease in BDI						38.6	.043
5b. % showing .33 <i>SD</i> decrease in mediator before .33 <i>SD</i> in outcome for the control condition Decrease in loneliness before decrease in BDI						15.5	<.001
Increase in emotional expression before decrease in BDI						10.7	<.001

Note. *B* = unstandardized regression coefficient; *SE* = standard error; Pr = partial regression coefficient;  $\Delta$  = change; BDI = Beck Depression Inventory; | = controlling for; *SD* = standard deviation.

depression. Both of these percentages were significantly lower than would be expected by chance. Although 72% of participants reported reading at least some of the book, it may take teens longer than 6 weeks to read this book and show symptom reduction. Indeed, bibliotherapy participants did show significantly greater reductions in depressive symptoms than did controls by 6-month follow-up (Stice et al., 2008).

Results provided only partial support for the intervention theory for the supportive expressive program. Relative to controls, supportive expressive participants did show clinically significant reductions in depressive symptoms and greater change in one mediator (emotional expression) but not the other mediator (loneliness). It is ironic that change in loneliness, but not in emotional expression, correlated with change in depression. Although the intervention effect on depression became nonsignificant when controlling for change in both of the mediators, the change in the main effect was significantly different only when controlling for change in loneliness. Data imply that only 15% showed a decrease in loneliness before showing a reduction in depression and that only 39% showed an increase in emotional expression before showing a reduction in depression (both percentages were significantly less than expected on the basis of chance).

There was mixed support for the specificity of the mediators. Group CB participants showed significant reductions in loneliness

and improvements in emotional expression, in addition to the expected effects for negative cognitions and pleasant activities, which suggests that this program affects both specific and non-specific factors. Supportive expressive group participants showed increases in pleasant activities but no change in negative cognitions. The former finding might suggest that reducing depression increases involvement in pleasant activities. Bibliotherapy did not reduce loneliness or emotional expression.

It is important to note that this five-step approach of testing mediators is not so stringent that it always fails to support intervention theory. In the first paper to apply this approach (Stice et al., 2007), we found that all five criteria were satisfied for a dissonance-based eating disorder prevention program. Indeed, the four other programs for which we have applied this test of mediation each failed for distinct reasons. This implies that this approach for testing mediation is sensitive to detecting violations of any of the five criteria implied by the mediational model.

Because change in depressive symptoms typically occurred before change in the mediator for all three prevention programs examined herein, it is important to consider alternative explanations for these findings. First, it is possible that the use of shortened versions of the mediator measures, but not the outcome, reduced the sensitivity of the former to detect intervention effects relative to the latter. Yet, the average pre- to postintervention effect (semipartial *r*s) was .17 for the mediators and .18 for the

BDI, and the average 3-week test–retest reliability among controls was similar for the mediators ( $r = .69$ ) and the BDI ( $r = .76$ ). Further, Stice et al. (2007) found full support for the mediational model of an eating disorder prevention program using an even briefer measure to assess the mediator.

Second, it may have been difficult to model the precise timing of change in this trial because we assessed the mediators and outcome only three times during the intervention. Although we could have modeled the timing of change more precisely with more frequent assessments, this factor cannot logically explain why results suggest that change in depression occurred before change in the mediators significantly more frequently than expected by chance for all three programs, as both mediators and the outcome were assessed three times. Further, if the estimated timing of changes was unreliable, one would expect to see only about half the sample showing changes in depression before showing changes in the mediators (the null hypothesis).

Third, it is possible that using slopes to test the temporal order of change in the mediators relative to the outcome is not optimal and that an alternative approach would provide evidence that the mediators change before symptoms. Taking an alternative approach, we calculated the proportion of participants in each condition who showed greater change in the mediator than in depression by the midintervention assessment. In the CB condition, the proportion of participants showing greater change in the mediator before change in the outcome in the expected direction was 17% for negative cognitions and 19% for pleasant activities. In the bibliotherapy condition, the proportion of participants showing greater change in the mediator before change in the outcome was 13% for negative cognitions and 14% for pleasant activities. In the supportive expressive condition, the proportion of participants showing greater change in the mediator before change in the outcome was 7% for emotional expression and 14% for loneliness. The fact that both approaches for modeling the relative timing of change produced similar results, despite different assumptions, implies the findings are robust.

Fourth, it is possible that regression to the mean resulted in more rapid reductions in depression than in the mediators, given that participants were initially selected because of elevated depression. However, the average effect for depression ( $r = .18$ ) was similar to the average effect for the mediators ( $r = .17$ ), which suggests that this alternative explanation is implausible.

The finding that change in depressive symptoms occurred more rapidly than change in all of the mediators may imply that nonspecific factors—such as the installation of hope; normalization of the person's experience; the creation of a therapeutic alliance, motivation, and a willingness to change; and the introduction to a theoretical model for change—drove symptom reductions. Yet, if nonspecific factors alone produced rapid reductions in depressive symptoms, it is difficult to explain why the group CB intervention outperformed the other interventions. From pre- to postintervention, CB group participants showed significantly greater reductions in BDI scores than did supportive expressive and bibliotherapy participants (Stice et al., 2008). Perhaps the group CB intervention was more credible and produced stronger nonspecific effects. Further, the CB intervention was the only program that resulted in significant improvements in all four mediators. This fact may suggest that

the effects were larger for this program because there were more mediators or because different mediators were operating for different subsets of participants. Because depression reductions often occur early, future trials testing for mediators of intervention effects should assess nonspecific factors in the first few sessions, as they may play an important role in generating intervention effects. This study has several limitations, including the reliance on self-report data, the infrequent measurement of mediators and the outcome, and the fact that we did not assess nonspecific factors. Future trials should investigate additional mediators that may account for depression prevention effects (e.g., nonspecific factors) and should attempt to assess mediators and the outcome more frequently during the intervention. They should also extend these procedures for testing mediational processes over longer follow-up periods, during which change may not be linear. Continued research on the mediators that produce intervention effects may result in more efficacious prevention programs.

## References

- Beck, A. T., Steer, R. M., & Garbin, M. (1988). Psychometric properties of the Beck Depression Inventory: 25 years of evaluation. *Clinical Psychology Review, 8*, 77–100.
- Hollon, S. D., & Kendall, P. C. (1980). Cognitive self-statements in depression: Development of an automatic thoughts questionnaire. *Cognitive Therapy and Research, 4*, 383–395.
- Jaycox, L. H., Reivich, K. J., Gillham, J., & Seligman, M. E. P. (1994). Prevention of depressive symptoms in school children. *Behaviour Research and Therapy, 32*, 801–816.
- Jolly, J. B., & Wiesner, D. C. (1996). Psychometric properties of the Automatic Thoughts Questionnaire with inpatient adolescents. *Cognitive Therapy and Research, 20*, 481–498.
- Kazdin, A. E., & Nock, M. K. (2003). Delineating mechanisms of change in child and adolescent therapy: Methodological issues and research recommendations. *Journal of Child Psychology and Psychiatry, 44*, 1116–1129.
- Lewinsohn, P. M., Roberts, R. E., Seeley, J. R., Rohde, P., Gotlib, I. H., & Hops, H. (1994). Adolescent psychopathology: II. Psychosocial risk factors for depression. *Journal of Abnormal Psychology, 103*, 302–315.
- MacPhillamy, D. J., & Lewinsohn, P. M. (1982). The Pleasant Events Schedule: Studies on reliability, validity, and scale intercorrelation. *Journal of Consulting and Clinical Psychology, 50*, 363–380.
- Rohde, P., Clarke, G. N., Mace, D. E., Jorgensen, J. S., & Seeley, J. R. (2004). An efficacy/effectiveness study of cognitive-behavioral treatment for adolescents with comorbid major depression and conduct disorder. *Journal of the American Academy of Child & Adolescent Psychiatry, 43*, 660–668.
- Rosenberg, M. (1979). *Conceiving the self*. New York, NY: Basic Books.
- Russell, D. W. (1996). UCLA Loneliness Scale (Version 3): Reliability, validity, and factor structure. *Journal of Personality Assessment, 66*, 20–40.
- Stice, E., Presnell, K., Gau, J., & Shaw, H. (2007). Testing mediators of intervention effects in randomized controlled trials: An evaluation of two eating disorder prevention programs. *Journal of Consulting and Clinical Psychology, 75*, 20–32.
- Stice, E., Rohde, P., Seeley, J., & Gau, J. (2008). Brief cognitive-behavioral depression prevention program for high-risk adolescents outperforms two alternative interventions: A randomized efficacy trial. *Journal of Consulting and Clinical Psychology, 76*, 595–606.
- Stice, E., Shaw, H., Bohon, C., Marti, C. N., & Rohde, P. (2009). A meta-analytic review of depression prevention programs for children and

- adolescents: Factors that predict magnitude of intervention effects. *Journal of Consulting and Clinical Psychology*, 77, 486–503.
- Weersing, V. R., & Weisz, J. R. (2002). Mechanisms of action in youth psychotherapy. *Journal of Child Psychology and Psychiatry*, 43, 3–29.
- Weissman, M. M., Orvaschel, H., & Padian, N. (1980). Children's symptom and social functioning self-report scales: Comparison of mothers' and children's reports. *Journal of Nervous and Mental Disease*, 168, 736–740.
- Yu, D. L., & Seligman, M. E. (2002). Prevention depressive symptoms in Chinese children. *Prevention & Treatment*, 5, Article 9. Available from <http://www.journals.apa.org/prevention/volume5/pre0050009a.html>

Received February 10, 2009

Revision received October 7, 2009

Accepted November 4, 2009 ■